

Strongly Asymmetric Tricriticality of Quenched Random-Field Systems

Alkan Kabakçioğlu^{1,2} and A. Nihat Berker^{1,2,3}

¹*Department of Physics, Massachusetts Institute of Technology
Cambridge, Massachusetts 02139, U.S.A.*

²*Feza Gürsey Research Center for Basic Sciences
Çengelköy, Istanbul 81220, TURKEY*

³*Department of Physics, Istanbul Technical University
Maslak, Istanbul 80626, TURKEY*

In view of the recently seen dramatic effect of quenched random bonds on tricritical systems, we have conducted a renormalization-group study on the effect of quenched random fields on the tricritical phase diagram of the spin-1 Ising model in $d = 3$. We find that random fields convert first-order phase transitions into second-order, in fact more effectively than random bonds. The coexistence region is extremely flat, attesting to an unusually small tricritical exponent β_u ; moreover, an extreme asymmetry of the phase diagram is very striking. To accomodate this asymmetry, the second-order boundary exhibits reentrance.

Tricritical phase diagrams of three-dimensional ($d = 3$) systems are strongly affected by quenched bond randomness: The first-order phase transitions are replaced, gradually as randomness is increased, by second-order phase transitions. The intervening random-bond tricritical point moves towards, and eventually reaches, zero-temperature, as the amount of randomness is increased. This behavior is an illustration of the general prediction that first-order phase transitions are converted to second-order by bond randomness [1–7], in a thresholded manner in $d = 3$. The randomness threshold increases from zero at the non-random tricritical point. The random-bond tricritical point maps, under renormalization-group transformations, onto a doubly unstable fixed distribution at strong coupling. Random-bond tricritical points exhibit a remarkably small value for the tricritical exponent $\beta_u = 0.02$, reflected in the near-flat top of the coexistence region. In the conversion of the first-order phase transition to second-order, traced by the random-bond tricritical point, a strong violation of the empirical universality principle occurs, via a renormalization-group fixed-point mechanism. Thus, detailed information now exists on the effect of quenched *bond* randomness on tricritical points, revealing several qualitatively distinctive features. [5,8]

No such information has existed on the effect of quenched *field* randomness on tricritical points. This topic is of interest also because renormalization-group studies have shown that quenched field randomness induces, under scale change, quenched bond randomness, as the presence of quenched field randomness continues. [9] Accordingly, we have conducted a global renormalization-group study of a tricritical system in $d = 3$ under quenched random fields. The results, presented below, show that these systems have their own distinctive behavior which is qualitatively different from that of non-random or random-bond tricritical systems. The latter distinction yields a microscopic physical intuition on the different effects of the two types of quenched

randomness.

We have studied the Blume-Emery-Griffiths (i.e., spin-1 Ising) model under quenched field randomness. The Hamiltonian is

$$-\beta\mathcal{H} = \sum_{\langle ij \rangle} \left[J s_i s_j + K s_i^2 s_j^2 - \Delta (s_i^2 + s_j^2) + H_{ij} (s_i + s_j) + H_{ij}^\dagger (s_i - s_j) \right], \quad (1)$$

where $s_i = \pm 1, 0$ at each site i of a simple cubic ($d = 3$) lattice and $\langle ij \rangle$ indicates summation over all nearest-neighbor pairs of sites. The quenched random fields H_{ij}, H_{ij}^\dagger are taken from a distribution

$$P(H, H^\dagger) = \frac{1}{4} \left[\delta(H + \sigma_H) \delta(H^\dagger + \sigma_H) + \delta(H + \sigma_H) \delta(H^\dagger - \sigma_H) + \delta(H - \sigma_H) \delta(H^\dagger + \sigma_H) + \delta(H - \sigma_H) \delta(H^\dagger - \sigma_H) \right]. \quad (2)$$

All other interactions in the initial Hamiltonian (1) are non-random. Under renormalization-group transformations, the Hamiltonian (1) maps onto a random-field random-bond Hamiltonian,

$$-\beta\mathcal{H} = \sum_{\langle ij \rangle} \left[J_{ij} s_i s_j + K_{ij} s_i^2 s_j^2 - \Delta_{ij} (s_i^2 + s_j^2) - \Delta_{ij}^\dagger (s_i^2 - s_j^2) + L_{ij} (s_i^2 s_j + s_i s_j^2) + L_{ij}^\dagger (s_i^2 s_j - s_i s_j^2) + H_{ij} (s_i + s_j) + H_{ij}^\dagger (s_i - s_j) \right], \quad (3)$$

where all interactions are quenched random, with a distribution function $P(J_{ij}, K_{ij}, \Delta_{ij}, \Delta_{ij}^\dagger, L_{ij}, L_{ij}^\dagger, H_{ij}, H_{ij}^\dagger)$ determined by the renormalization-group transformation. Specifically, the first four arguments here reflect the rescaling-induced bond randomness of the random-field system.

The renormalization-group transformation is contained in the mapping between the quenched probability distributions of the starting and rescaled systems,

$$P'(\vec{K}'_{ij'}) = \int \left[\prod_{\langle ij \rangle} d\vec{K}_{ij} P(\vec{K}_{ij}) \right] \delta(\vec{K}'_{ij'} - \vec{R}(\{\vec{K}_{ij}\})), \quad (4)$$

where $\vec{K}_{ij} \equiv (J_{ij}, K_{ij}, \Delta_{ij}, \Delta_{ij}^\dagger, L_{ij}, L_{ij}^\dagger, H_{ij}, H_{ij}^\dagger)$ are the interactions at locality $\langle ij \rangle$, the primes refer to the renormalized system, the product is over all unrenormalized localities $\langle ij \rangle$ whose interactions $\{\vec{K}_{ij}\}$ influence the renormalized interaction $\vec{K}'_{ij'}$, and $\vec{R}(\{\vec{K}_{ij}\})$ is a local recursion relation that embodies the latter dependence. Simply said, Eq.(4) sums over the joint probabilities of the values of neighboring unrenormalized interactions that conspire to yield a given value of the renormalized interaction. The phenomena characteristic to quenched randomness should derive from the probability convolution shown in Eq.(4), rather than the precise form of the recursion \vec{R} that should be a smooth local function. In this work, we use the Migdal-Kadanoff recursion

relation, given for this system in [10]. The convolution is effected by representing $P(\vec{K}_{ij})$ in terms of bins, the degree of detail (i.e., the number of bins) reflecting the level of approximation. In this work, we have used 531,441 bins, corresponding to renormalization-group flows in a 4,782,969-dimensional space. The application of Eq.(4) via the binning procedure has been described elsewhere [8,11].

Our main result, the striking difference between the three types of $d = 3$ Ising tricriticality, is evident in Fig.1, where the calculated random-field, random-bond, and non-random phase diagrams are superimposed. The same amount of quenched randomness [$\sigma_H = 0.2 = \sigma_\Delta$, as in Eq.(2)] is used, for relevance of comparison. It is seen that both bond randomness and field randomness convert first-order phase transitions to second-order in a thresholded manner, but that field randomness is more effective than bond randomness in this conversion. Both types of random tricritical points occur at remarkably near-flat tops of coexistence regions, reflecting the unusually small values of the exponent β_u , but the random-field phase diagram is most strikingly asymmetrical. The tricritical point occurs at the high density $\langle s_i^2 \rangle = 0.835$ (as opposed to 0.613 and 0.665 in the random-bond and non-random systems, respectively), essentially all of the near-flat portion of the coexistence top occurring on the dilute branch of the coexistence boundary. To accommodate this asymmetry, the randomness-extended line of second-order phase transitions has to curve over and exhibit reentrant behavior [12] as a function of temperature. This difference in behavior comes from the fact that random bonds destroy order-disorder coexistence without destroying order itself [3,4], whereas random fields destroy both order-disorder coexistence, through the rescale-generated bond randomness, and order per se. The latter is more effective near the tricritical point, where considerable vacancy fluctuations occur within the ordered phase.

Random-field and random-bond tricritical points renormalize onto obviously different doubly unstable fixed distributions (the field variables L and H remain at zero in the latter case).

It is seen in Fig.1 that the coexistence boundary of either type of random system follows that of the non-random system, until the temperature-lowered tricritical region sets in relatively abruptly. This is similar to the magnetization of random-field systems following the non-random curve until the critical region sets in quite abruptly. [13,11]

On the high-temperature side of the tricritical point, it has been known [8] that the break in slope of the critical line, in the random-bond system, is connected to the strong violation of the empirical universality principle, segments on each side of this point having their own critical exponents, respectively of the strong-coupling and non-strong-coupling type. No such universality violation occurs along the second-order line of the random-field

system, the entire line having random-field critical exponents that are governed by a strong-coupling fixed distribution, implying a modified hyperscaling relation [14,15].

The global phase diagrams of the random-field and random-bond systems are given in Figs.2(a,b). It is seen that all first-order phase transitions are completely converted to second-order, at a zero-temperature tricritical point, for $\sigma_H \simeq 0.5$ and $\sigma_\Delta \simeq 0.7$, respectively. Furthermore, the ordered phase is eliminated at $\sigma_H \simeq 0.9$ in the random-field case, but persists for all for σ_Δ in the random-bond case.

This research was supported the U.S. Department of Energy under Grant No. DE-FG02-92ER45473 and by the Scientific and Technical Research Council of Turkey (TÜBİTAK). We gratefully acknowledge the hospitality of the Feza Gürsey Research Center for Basic Sciences and of the Istanbul Technical University.

-
- [1] J. Wehr and M. Aizenman, Phys. Rev. Lett. **62**, 2503 (1989).
 - [2] K. Hui and A.N. Berker, Phys. Rev. Lett. **62**, 2507 (1989); **63**, 2433(E) (1989).
 - [3] A.N. Berker, J. Appl. Phys. **70**, 5941 (1991).
 - [4] A.N. Berker, Physica A **194**, 72 (1993).
 - [5] A.N. Berker and A. Falicov, Turk. J. Phys. **18**, 347 (1994).
 - [6] S. Chen, A.M. Ferrenberg, and D.P. Landau, Phys. Rev. Lett. **69**, 1213 (1992).
 - [7] S. Chen, A.M. Ferrenberg, and D.P. Landau, Phys. Rev. E **52**, 1377 (1995).
 - [8] A. Falicov and A.N. Berker, Phys. Rev. Lett. **76**, 4380 (1996).
 - [9] S.R. McKay and A.N. Berker, J. Appl. Phys. **64**, 5785 (1988).
 - [10] A. Lopatnikova and A.N. Berker, Phys. Rev. B **56**, 11865 (1997).
 - [11] A. Falicov, A.N. Berker, and S.R. McKay, Phys. Rev. B **51**, 8266 (1995).
 - [12] P. Cladis, Mol. Cryst. Liq. Cryst. **165**, 85 (1988).
 - [13] M.S. Cao and J. Machta, Phys. Rev. B **48**, 3177 (1993).
 - [14] A.J. Bray and M.A. Moore, J. Phys. C **18**, L927 (1985).
 - [15] A.N. Berker and S.R. McKay, Phys. Rev. B **33**, 4712 (1986).

Figure Captions

Fig. 1: Calculated tricritical phase diagrams for non-random (dotted), random-bond (full), and random-field (dashed) $d = 3$ systems for $K = 0$. In each phase diagram, a line of second-order phase transitions extending to high temperatures meets, at a tricritical point, a coexistence region extending to low temperatures. Note the

near-flat top of the coexistence regions in both quenched random systems, and the extreme asymmetry of the random-field system. Thus, field randomness is more effective than bond randomness in converting first-order transitions into second-order (i.e., the tricritical point is at lower temperature).

Figs 2: Calculated $d = 3$ global phase diagrams for $K = 0$: (a) Random-bond systems exhibit two universality classes of second-order phase transitions (thin and thick full lines) and first-order phase transitions (circles). (b) Random-field systems exhibit second-order (full lines) and first-order (circles) phase transitions. In both types of quenched randomness, the first-order transitions cede under increased randomness. The line of tricritical points (dashed) reaches zero temperature, as all transitions become second-order. In the random-field case, the ordered phase disappears under further randomness, whereas in the random-bond case, the ordered phase (and the strong violation of universality) persists for all randomness.




FIG.1




FIG.2a




FIG.2b